

Student:	Grade:	School Contact	-•	DOB:
Asthmatic: 🗖 Yes 📮	No (increased risk for seve	ere reaction) Severity of 1	reaction(s):	
Mother:		MHome #:	MWork #:	MCell #:
Father:		FHome #:	FWork #:	FCell #:
Emergency Contact:		Relationship:	p	hone:
 MOUTH THROAT SKIN STOMACH LUNG HEART The 	ALLERGIC REACTION Itching & swelling of lips Itching, tightness in throw Hives, itchy rash, swelling Nausea, abdominal cramy Shortness of breath, repe "Thready pulse", "passin e severity of symptoms important that treatmo	, tongue or mouth at, hoarseness, cough g of face and extremities ps, vomiting, diarrhea titive cough, wheezing g out" s can change quickly	_	SE: Student Photo
STAFF MEMBERS IN		Classroom Teacher(s)Support Staff	1	Area Teacher(s) ortation Staff
TREATMENT: Remove stinger if visible, apply ice to area. Rinse contact area with water. Treatment should be initiated with symptoms without waiting for symptoms Benadryl ordered: Yes No Give				
Call school nurse. Call parent/guardian if off school grounds.				
Epinephrine ordered:	🛛 Yes 📮 No	Special instructions:		
IF ANY SYMPTOMS BEYOND REDNESS OR SWELLING AT THE SITE OF THE STING ARE PRESENT AND EPINEPHRINE IS ORDERED, GIVE EPINEPHRINE IMMEDIATELY AND CALL 911. Preferred Hospital if transported: Epinephrine provides a 20 minute response window. After epinephrine, a student may feel dizzy or have an increased heart rate. This is a normal response. Students receiving epinephrine should be transported to the hospital by ambulance. A staff member should accompany the student to the emergency room if the parent, guardian or emergency contact is not present and adequate supervision for other students is present.				
1	☐ Medication available on			
Healthcare Provider:			Phone:	
Written by:				
Written by:				

This plan is in effect for the current school year and summer school as needed.